

COMMENTARY

On the Performance of Continuum Solvation Methods. A Comment on “Universal Approaches to Solvation Modeling”

CON SPECTUS

In a recent Account, Cramer and Truhlar presented a comparison between the SM8 method and standard versions of other continuum solvation models implemented in widely available quantum mechanical programs. In that Account, the SM8 model was found to lead to “considerably smaller errors for aqueous and nonaqueous free energies of solvation for neutrals, cations, and anions, with particularly good performance for nonaqueous data”. Here, we demonstrate that competing solvation methods are indeed as accurate as the SM8 method, if they are applied with the same rigor.

In their recent Account,¹ “A Universal Approach to Solvation Modeling”, Cramer and Truhlar provide a concise description of the performance of the SM8 solvation model, which has been described in detail in a recent paper.² The SM8 model is the latest version of the series of SM_x continuum methods and has been parametrized using the experimental solvation free energies of a large set of compounds containing a variety of functional groups in different solvents. These data, which consist of 2489 entries for 940 neutral compounds and 332 ionic ones, have been used in conjunction with the solvent radius parametrized previously to optimize a large number of empirical model parameters, including solvent-dependent Coulomb radii, atomic surface tensions, and other parameters contained in the charge models and in the generalized Born model. The SM8 model is described as a continuum solvation model universally applicable to general solvents, for which a small number of physicochemical descriptors are known either experimentally or from estimation.

In Table 1 of their Account, the authors demonstrate that the SM8 model succeeds in reproducing the solvation free energies of the training data set with a mean unsigned error (MUE) of 0.59 kcal/mol for neutral solutes and 4.31 kcal/mol for ions. A comparison with four other solvation models is also presented. One of them is the Poisson–Boltzmann method implemented in the Jaguar program^{3,4} and the other three are (1) the integral equation formalism of the polarizable continuum model, IEF-PCM,⁵ which is the default version of the PCM family of methods⁶ as imple-

mented in the Gaussian03⁷ package, (2) the C-PCM method,⁸ which is an implementation of the conductor-like screening ansatz COSMO⁹ in the framework of the PCM model (the most effective version of the method is also available in Gaussian03¹⁰), and (3) the COSMO implementation¹¹ in the NWCHEM program,¹² which is incorrectly referenced as GCOSMO by Cramer and Truhlar.^{1,2}

For the same set of training compounds, MUEs of 5.66, 2.43, and 4.29 kcal/mol are reported for the solvation free energies predicted for neutral solutes (and even larger MUEs for ionic species) using the self-consistent reaction field (SCRF) continuum solvation models IEF-PCM, C-PCM, and NWCHEM/COSMO, respectively. We consider that the conclusions that a nonexpert reader can obtain might be inappropriate, since they are based on the comparison of the solvation free energies predicted by using the highly trained SM8 model with the IEF-PCM, C-PCM, and COSMO models implemented in widely available programs, which were not trained in the same way. Other more elaborate versions providing higher accuracy for the computation of solvation free energies have been reported elsewhere (see below). This point must be stressed in order to avoid a wrong impression about the performance of “non-SM_x models” for non-expert users.

In order to examine the ability of IEF-PCM, C-PCM, and COSMO models to estimate solvation free energies, we have determined the solvation free energy with the IEF-PCM and COSMO-RS^{13,14} models using the MST (Miertus–Scrocco–Tomasi)^{15,16} and COSMOtherm_C2.1_0107¹⁷ versions, respectively, for subsets of

TABLE 1. Statistical Performance (Mean Unsigned Error in kcal/mol) of COSMOtherm and SM8 on Solvation Free Energies of Neutral Compounds

data set	<i>N</i>	COSMOtherm	SM8
all neutral	2346	0.48	0.59
subset of 17 solvents	960	0.56	0.61
aqueous solvation	284	0.58	0.55

neutral compounds included in the SM8 training set. These versions represent in our view the state-of-the-art for these methods when applied to the calculation of solvation free energies (due to the close similarity between IEF-PCM and C-PCM results for a common set of parameters, if C-PCM is used with the scaling factor $(\epsilon-1)/(\epsilon+0.5)$, the latter method will not be discussed here).

With this aim, we retrieved the chemical structures of all solutes from the chemical names given in the Supporting Information of the original SM8 paper.² Then, we used the standard procedures established for IEF-PCM/MST and COSMO-RS to determine the solvation free energies. In essence, IEF-PCM/MST computations were performed at both HF and DFT (B3LYP) levels of theory using the 6-31G(d) basis and molecular geometries optimized in the gas phase, whereas COSMO-RS results were obtained at the DFT level (BP functional with TZVP basis set) and geometries separately optimized in vacuum and at $\epsilon = \infty$. Multiple conformations were taken into account self-consistently where necessary. Computations were performed using Gaussian03 for IEF-PCM/MST and TURBOMOLE¹⁸ and COSMOtherm programs for COSMO-RS, respectively, with no additional fitting of any parameters.

The statistical performance of COSMO-RS can be examined from the results given in Table 1, which also reports the corresponding SM8 results taken from Table 1 in the reference paper.¹ Since COSMOtherm does not require individual quantum chemical calculations of each solute in each solvent but only one per solute in a perfect conductor, COSMOtherm was applied to all 91 solvents and 2346 solute/solvent pairs given in the Supporting Information of the SM8 paper. The results show that the performance of COSMO-RS is identical to or slightly better than that of the SM8 method, even for the compounds used in the SM8 training set. It must be noted that no experimental or adjustable solvent parameters have been used in COSMOtherm calculations, because the solvents are described on the same DFT/COSMO basis as the solutes within the COSMO-RS theory. The largest deviations in COSMOtherm calculations are found for thiophosphate compounds, for which the structures are slightly questionable and the experimental

TABLE 2. Statistical Performance (Mean Unsigned Error in kcal/mol) of IEFPCM-MST and SM8 on Solvation Free Energies of Neutral Compounds

data set	<i>N</i>	IEF-PCM/MST HF	IEF-PCM/MST B3LYP	SM8
nonaqueous solvation	960 ^a	0.64	0.64	0.61
aqueous solvation	284 ^b	1.01	1.01	0.55

^a IEF-PCM/MST calculations performed on 350 compounds and 3 solvents (octanol, CHCl₃, CCl₄). ^b IEF-PCM/MST calculations performed on 237 compounds.

data show at least some strange trends. The straight applicability and good performance of COSMO-RS for all 91 solvents also proves that COSMO-RS is a universal solvation model. Indeed, the "universal" nature of COSMO-RS is further underlined by its ability to treat mixtures of solvents without the need for any experimental solvent data.

Since the MST version of the IEF-PCM has been parametrized for water, octanol, chloroform, and carbon tetrachloride, the statistical analysis reported in Table 2 is limited to the subset of experimental data available for these solvents. In addition, compounds containing P, Br, and Si were eliminated from the SM8 data set because these elements were not considered in the MST parametrization.

The results show that the MUEs of IEF-PCM/MST and SM8 results are nearly the same for compounds in octanol, chloroform, and carbon tetrachloride. Moreover, the MUE resembles that obtained in the original parametrization of the IEF-PCM/MST method for each solvent using a smaller set of compounds (root mean square deviation, rmsd, of 0.6/0.6, 0.5/0.5, and 0.3/0.3 kcal/mol in octanol, chloroform, and carbon tetrachloride at the HF/B3LYP level for 63, 56, and 48 compounds).^{15,16} A larger MUE is found in water, also slightly larger than that obtained in the original MST parametrization (rmsd of 0.8/0.8 for 72 compounds). This trend is not surprising and can be mainly attributed to the limited number of optimized empirical parameters used in the MST implementation to maintain the general applicability of the IEF-PCM formalism and avoid overtraining.

Overall, the preceding discussion allows us to conclude that IEF-PCM, C-PCM, and COSMO-RS methods are capable of predicting the experimental solvation free energies of (bio)organic compounds in a variety of solvents. The only premise is that the user must keep the specific details of the computational protocol adopted in the parametrization of each continuum model in order to maintain the balance between electrostatic and nonelectrostatic components.¹⁹ In particular, the default implementations of IEF-PCM (and C-PCM) in Gaussian03, which have been used in the reference paper,¹ have never been quantitatively parametrized

for the reproduction of solvation free energies. As a result, the default set of radii (known with the UA0 acronym) used to build up the solute/solvent interface is not the best choice for the quantitative prediction of solvation free energies. As a matter of fact, such a cavity represents a compromise to gain qualitative insight into the role played by solvation on a variety of properties of the solute without the need to change the cavity parameters each time. At this point, we should emphasize that the accurate prediction of solvation free energies is not the only requirement for a solvation method, which must also reliably predict solvent effects on geometrical, vibrational, electronic and magnetic properties, as well as on many response properties of any solvated system.^{6,20,21}

Conclusions

Understanding of the solvent effect on the properties of molecules and chemical processes still represents a challenge for the community of theoretical chemists, which explains the variety of computational strategies developed for the study of chemistry in condensed phases. Among these strategies, SCRF methods, despite the rather crude approximation of representing the solvent as a polarizable continuum medium, have a number of advantages, such as the definition of computational algorithms based on well established physical laws, the simplicity of the mathematical formalism, and the reduced computational cost. The progress made in the last decades in SCRF methods has crystallized in a series of optimized continuum methods, like (among others) SM8, PCM (either IEF-PCM or C-PCM) or COSMO-RS, which yield accurate solvation free energies provided that the user of these methods pays attention to the specific details of the parametrization made for each model.

This is an encouraging achievement that should stimulate further studies aimed at the calibration of the most advanced implementations of continuum models for understanding the physical and chemical properties of solutes in solution. In this context, we think that the concept of “universal” solvation models should not be limited exclusively to the ability to estimate reliably the solvation free energy of chemical species in a large number of solvents, but also to the most fundamental problem of providing useful information about the response of the solute to the condensed phase environment and its influence on a variety of physical and chemical properties.

Andreas Klamt,^{*,†‡} Benedetta Mennucci,[§] Jacopo Tomasi,[§] Vincenzo Barone,[†] Carles Curutchet,^{||} Modesto Orozco,^{||} and F. Javier Luque[▽]

[†]*COSMOlogic GmbH&CO KG, Burscheider Str. 515, 51381 Leverkusen, Germany*, [‡]*Department of Physical and Theoretical Chemistry, University of Regensburg, 93040 Regensburg, Germany*, [§]*Department of Chemistry, University of Pisa, Via Risorgimento 35, 56126 Pisa, Italy*, [†]*Scuola Normale Superiore, piazza dei Cavalieri 7, 56126, Pisa* and *IPCF-CNR, via Moruzzi 1, 56100 Pisa, Italy*, ^{||}*Department of Chemistry, University of Toronto, 80 St. George Street, Toronto, Ontario, M5S 3H6 Canada*, ^{||}*Molecular Modeling and Bioinformatics Unit, Institut de Recerca Biomèdica. Barcelona Scientific Park, Josep Samitier 1–6, 08028 Barcelona, Spain*, *Department of Life Sciences, Barcelona Supercomputing Centre, Jordi Girona 29, 08034 Barcelona, Spain*, and *Departament de Bioquímica, Facultat de Biologia, Avda Diagonal 647, Barcelona 08028, Spain*, [▽]*Department of Physical Chemistry and Institute of Biomedicine (IBUB), Faculty of Pharmacy, University of Barcelona, Avda. Diagonal 643, 08028 Barcelona, Spain*

FOOTNOTES

*Corresponding author. Tel: +49-2171-731681. E-mail: klamt@cosmologic.de.

REFERENCES

- Cramer, C. J.; Truhlar, D. G. A Universal Approach to Solvation Modeling. *Acc. Chem. Res.* **2008**, *41*, 760–768.
- Marenich, A. V.; Olson, R. M.; Kelly, C. P.; Cramer, C. J.; Truhlar, D. G. Self-Consistent Reaction Field Model for Aqueous and Nonaqueous Solutions Based on Accurate Polarized Partial Charges. *J. Chem. Theory Comput.* **2007**, *3*, 2011–2033.
- Tannor, D. J.; Marten, B.; Murphy, R.; Friesner, R. A.; Sitkoff, D.; Nicholls, A.; Ringnalda, M.; Goddard, W. A.; Honig, B. Accurate First Principles Calculation of Molecular Charge Distributions and Solvation Energies from Ab Initio Quantum Mechanics and Continuum Dielectric Theory. *J. Am. Chem. Soc.* **1994**, *116*, 11875–11882.
- Marten, B.; Kim, K.; Cortis, C.; Friesner, R. A.; Murphy, R. B.; Ringnalda, M. N.; Sitkoff, D.; Honig, B. New Model for Calculation of Solvation Free Energies: Correction of Self-Consistent Reaction Field Continuum Dielectric Theory for Short-Range Hydrogen-Bonding Effects. *J. Phys. Chem.* **1996**, *100*, 11775–11788.
- Cances, E.; Mennucci, B.; Tomasi, J. A New Integral Equation Formalism for the Polarizable Continuum Model: Theoretical Background and Applications to Isotropic and Anisotropic Dielectrics. *J. Chem. Phys.* **1997**, *107*, 3032–3041.
- Tomasi, J.; Mennucci, B.; Cammi, R. Quantum Mechanical Continuum Solvation Models. *Chem. Rev.* **2005**, *105*, 2999–3093.
- Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A., Jr.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. *Gaussian 03*, revision D.01; Gaussian, Inc.: Wallingford, CT, 2004.
- Barone, V.; Cossi, M. Quantum Calculation of Molecular Energies and Energy Gradients in Solution by a Conductor Solvent Model. *J. Phys. Chem. A* **1998**, *102*, 1995–2001.
- Klamt, A.; Schüürmann, G. COSMO: A New Approach to Dielectric Screening in Solvents with Explicit Expressions for the Screening Energy and its Gradient. *J. Chem. Soc., Perkin Trans.* **1993**, *2*, 799–805.

- 10 Cossi, M.; Rega, N.; Scalmani, G.; Barone, V. Energies, Structures, And Electronic Properties of Molecules in Solution with the C-PCM Solvation Model. *J. Comput. Chem.* **2003**, *24*, 669–681.
- 11 Liotard, D. A.; Hawkins, G. D.; Lynch, G. C.; Cramer, C. J.; Truhlar, D. G. Improved Methods for Semiempirical Solvation Models. *J. Comput. Chem.* **1995**, *16*, 422–440.
- 12 Bylaska, E. J.; et al. *NWChem, A Computational Chemistry Package for Parallel Computers*, version 4.7; Pacific Northwest National Laboratory: Richland, WA, 2006.
- 13 Klamt, A. Conductor-like Screening Model for Real Solvents: A New Approach to the Quantitative Calculation of Solvation Phenomena. *J. Phys. Chem.* **1995**, *99*, 2224–2235.
- 14 Klamt, A. *COSMO-RS from Quantum Chemistry to Fluid Phase Thermodynamics and Drug Design*; Elsevier: Amsterdam, 2005.
- 15 Curutchet, C.; Orozco, M.; Luque, F. J. Solvation in Octanol: Parametrization of the Continuum MST Model. *J. Comput. Chem.* **2001**, *22*, 1180–1193.
- 16 Soteras, I.; Curutchet, C.; Bidon-Chanal, A.; Orozco, M.; Luque, F. J. Extension of the MST Model to the IEF Formalism: HF and B3LYP Parametrizations. *J. Mol. Struct. (THEOCHEM)* **2005**, *727*, 29–40.
- 17 Eckert, F.; Klamt, A. *COSMOtherm*, Version C2.1, Revision 01.07; COSMOlogic GmbH&CoKG, Leverkusen, Germany 2007.
- 18 Turbomole, version 5.10, <http://www.turbomole.com>, 2008.
- 19 Curutchet, C.; Cramer, C. J.; Truhlar, D. G.; Ruiz Lopez, M.; Rinaldi, D.; Orozco, M.; Luque, F. J. Electrostatic Component of Solvation: Comparison of SCRF Continuum Models. *J. Comput. Chem.* **2003**, *24*, 284–297.
- 20 Barone, V.; Improta, R.; Rega, N. Quantum Mechanical Computations and Spectroscopy: From Small Rigid Molecules in the Gas Phase to Large Flexible Molecules in Solution. *Acc. Chem. Res.* **2008**, *41*, 605–616.
- 21 Mennucci, B., Cammi, R., Eds. *Continuum solvation models in chemical physics, from theory to applications*; Wiley: Chichester, U.K., 2007.